




- ◆ Identify, Isolate, Inform: Ambulatory Care Evaluation of Patients with Possible Ebola Virus Disease (Ebola)

WASHOE COUNTY HEALTH DISTRICT • P.O. BOX 11130 • RENO, NEVADA • 89520-0027 • (775) 328-2447

This alert contains information on investigations in progress and/or diagnoses that may not yet be confirmed.

The Physician Alert is intended primarily for the use of local health care providers, should be considered privileged, and should **NOT** be distributed further.

IDENTIFY, ISOLATE, INFORM: AMBULATORY CARE EVALUATION OF PATIENTS WITH POSSIBLE EBOLA VIRUS DISEASE (EBOLA)



On November 1, 2014, CDC issued a new algorithm entitled "Identify, Isolate, Inform: Ambulatory Care Evaluation of Patients with Possible Ebola Virus Disease (Ebola)" (see following page). This new algorithm specifically targets Ambulatory Care facilities. Washoe County Health District (WCHD) highly recommends all healthcare providers in ambulatory care settings follow this algorithm. Again, travel history to West African countries with widespread Ebola transmission (Guinea, Liberia, and Sierra Leone as of November 3, 2014) or an exposure history to suspect cases are critical components in the evaluation process. Should you have any patients who meet criteria for a Person Under Investigation (PUI) for Ebola, coordination efforts should be made among your facility, WCHD, and the receiving hospital. Should you have any questions regarding this evaluation algorithm, please call **WCHD at 775-328-2447**.

The telephone numbers for local emergency departments are as follows:

Renown 775-982-4144

Renown at South Meadows: 775-982-7144

St. Mary's Regional Medical Center 775-770-3173

Northern Nevada Medical Center 775-356-4040


Incline Village Community Hospital 775-888-4210

Please share this document with all physicians/staff in your office/facility.

Identify, Isolate, Inform: Ambulatory Care Evaluation of Patients with Possible Ebola Virus Disease (Ebola)



The majority of febrile patients in ambulatory settings do not have Ebola Virus Disease (Ebola), and the risk posed by Ebola patients with early, limited symptoms is lower than that from a patient hospitalized with severe disease. Nevertheless, because early Ebola symptoms are similar to those seen with other febrile illnesses, triage and evaluation processes should consider and systematically assess patients for the possibility of Ebola.



YES - Patient may meet criteria for Person Under Investigation for Ebola*

- 3 Isolate patient immediately: Avoid unnecessary direct contact**
- Place patient in private room or area, preferably enclosed with private bathroom or covered commode.
 - Avoid unnecessary direct contact.
 - If direct contact is necessary, personal protective equipment (PPE) and dedicated equipment must be used to minimize transmission risk.
 - Only essential personnel with designated roles should evaluate patient.
 - If patient is exhibiting obvious bleeding, vomiting or copious diarrhea, then do not re-enter room until EMS personnel trained to transport Person Under Investigation for Ebola arrive.
 - Do not perform phlebotomy or any other procedures unless urgently required for patient care or stabilization.
 - Consult with the health department before cleaning up blood or body fluids. Any reusable equipment should not be reused until it has been appropriately cleaned and disinfected.*

AND

- 4 Inform Health Department and prepare for safe transport.**
- Contact the relevant health department IMMEDIATELY.
 - Prepare for transfer to a hospital identified by the health department for evaluation of possible Ebola.
 - Coordinate with health department regarding:
 - Who will notify the receiving emergency department or hospital about the transfer, and
 - Arrangements for safe transport to accepting facility designated by public health officials.

PERSONS UNDER INVESTIGATION FOR EBOLA SHOULD ONLY BE SENT TO HOSPITALS AND FACILITIES SPECIFICALLY DESIGNATED BY PUBLIC HEALTH OFFICIALS.

Do not transfer without first notifying the health department.

PPE in the ambulatory care setting**:

- No one should have direct contact with a Person Under Investigation for Ebola without wearing appropriate personal protective equipment (PPE).
- If PPE is available and direct patient contact necessary, a single staff member (trained in proper donning and removal of PPE) should be designated to interact with the Person Under Investigation.
- At a minimum, health care workers should use the following PPE before direct patient contact:
 - A.** Face shield & surgical face mask,
 - B.** Impermeable gown, and
 - C.** Two pairs of gloves.
- The designated staff member should refrain from direct interaction with other staff and patients in the office until PPE has been safely removed in a designated, confined area. Examples of safe donning and removal of PPE should be reviewed: http://www.cdc.gov/hicpac/2007IP/2007ip_fig.html

NOTE: Patients with exposure history and Ebola-compatible symptoms seeking care by phone should be advised to remain in place, minimize exposure of body fluids to household members or others near them, and given the phone number to notify the health department. The ambulatory care facility must also inform the health department. If the clinical situation is an emergency, the ambulatory care facility or patient should call 911 and tell EMS personnel the patient's Ebola risk factors so they can arrive at the location with the correct PPE.

*Refer to <http://www.cdc.gov/vhf/ebola/> for the most up-to-date guidance on the **Case Definition for Ebola, Environmental Infection Control and Ebola-Associated Waste Management**;

**Refer to <http://www.cdc.gov/hai/settings/outpatient/outpatient-care-guidelines.html> for a summary guide of infection prevention recommendations for outpatient settings.